

Estimating PM_{2.5} in the Northeast United States Using Kriging


Shifali Reddy, Daniel Vidal, Nabin Malakar, Fred Moshary, Barry Gross

The City College
of New York


The City College of New York

Background and Objectives

- PM_{2.5} is a form of particle pollution and is found primarily in smoke and haze.
- Particle pollution contains microscopic solids or liquid droplets that are able to enter deep into the lungs and cause serious health problems.
- PM_{2.5} has been linked to increased respiratory symptoms and cardiovascular disorders.
- Our goals from the research project are to obtain kriging results with variance and to obtain fusion between remote sensing and kriging.



Approximate size of PM_{2.5}



PM_{2.5} Health Effects


Materials and Method

Data:

- AQS (Air Quality System, EPA) is a repository of ambient air quality data from over 10,000 EPA, state, local, and tribal pollution control agencies.
- RSIG (Remote Sensing Information Gateway, EPA) is a tool for selecting, subsetting, downloading, assembling, and visualizing environmental data. It renders each dataset and overlays them on a map.

Kriging

- Kriging, a method of interpolation, is based on regression against observed values of surrounding data points, weighted according to a covariance function.
- Simple kriging assumes the expectation of the random field to be known.
- If the data points are dense and uniformly distributed, we expect to obtain good results regardless of an interpolation algorithm.
- If the data points fall in a few clusters with large gaps in between, we expect to obtain unreliable estimates regardless of an interpolation algorithm.
- We expect best estimates when the PM_{2.5} values fall under a normal distribution.
- Kriging compensates for the effects of data clustering by putting more weight on isolated points and treating clusters like single points.
- The kriging algorithm will need to access to semivariogram values for lag distances other than those used in the empirical variogram.
- Semi variogram models used need to obey certain numerical properties for the kriging equations to be solvable.
- We used the spherical model, which exhibits linear behavior at the origin, appropriate for representing properties with a higher level of short-range variability.



Results




Fig. 1




Fig. 2




Fig. 3




Fig. 4









Fig. 5



May 24, 2008



September 25, 2006



Summary and Conclusion

- The box plot demonstrates how PM_{2.5} levels are elevated during the summer months of June, July and August.
- We implemented kriging, specifically simple kriging, to obtain better spatial coverage for the point station measurements.
- We also obtained the estimation as well as variance for the PM_{2.5} measurement for the northeastern region of the United States.
- Furthermore, these products will enable us to obtain a fused data product, fusing together ground station data and satellite remote sensing estimates.

Acknowledgements: The authors greatly appreciate the National Oceanic and Atmospheric Administration Cooperative Remote Sensing Science and Technology Center (NOAA CREST), the New York City Research Initiative (NYCRI), the Environmental Protection Agency (EPA), and the Goddard Institute for Space Studies (GISS) for their support. The authors also appreciate Pedro Placido for his tremendous help and education provided.